

AFTOVAXPUR
PURIFIED INACTIVATED VACCINE
AGAINST FOOT AND MOUTH DISEASE IN RUMINANTS

CATEGORY FOR DISTRIBUTION PP (Vet)

COMPOSITION

Antigen:

Purified Inactivated Foot and Mouth Disease Virus containing any type of 5 serotypes appropriate to the region. Type SAT 1, Type SAT 2, Type SAT 3, type O, Type A

Adjuvants:

Aluminium hydroxide and saponin

INDICATIONS

The vaccination of cattle, buffalo, sheep and goats against Foot and Mouth Disease

ADMINISTRATION AND DOSAGE

Shake well before use

Subcutaneous route

Dose: regardless of the age and weight of the animal. Refer to the label on the container according to the following schedule:

Primary course of vaccination:

Two injections, 3 to 4 weeks apart, starting at two weeks of age for young animals from unvaccinated dams or 2.5 months of age for young animals from vaccinated dams. In case of an epidemic the first injection should be given to all animals as early as 2 weeks of age.

Boosters:

Normally, boosters should be given every 4- 6 months depending on the risk and in accordance with local legislation.

CONTRADICTIONS:

None

PRECAUTIONS

Vaccinate only healthy animals

Observe usual aseptic procedures. Once opened bottles of vaccines should be used within 24 hours provided they have been stored between +2°C and 8°C.

SIDE EFFECTS

Vaccination may be followed by a small local swelling and/ or slight pyrexia both of short duration.

WITHDRAWAL TIME

Nil

STORAGE

Store between +2°C and 8°C, protected from the light. **Do not freeze.**

PRESENTATION

300 ml Bottle and 100ml Bottle

FOR VETERINARY USE

| | |
|--|--|
| Produced with Boehringer Ingelheim Animal Health Technology | Manufactured by BOTSWANA VACCINE INSTITUTE P/Bag 0031 Gaborone , Botswana Tel: +267 391 2711 Email: marketing@bvi.co.bw |
|--|--|

